


«Протез синовиальной жидкости Hyruan Inj. в шприце объемом 2,5 мл»
Протез синовиальной жидкости (1 шт. в упаковке)





LG Life Sciences

Prefilled type intra-articular injection

Hyruan Inj.


Sodium Hyaluronate 1%

NOT FOR SALE

Lot No.: HIR12005
Mfg. Date: 2012.06.15
Exp. Date: 2015.06.14

25 mL x 5 Syringes/pack

10 mg/mL



Viscosupplementation Treatment for Arthritic Joints

Hyruan Inj.

Sodium Hyaluronate 1%

used for patients with an osteoarthritis-related pain and stiffness in the affected joints, chronic, and gradual therapy. After the injection of Hyruan Inj., which is a natural product of the body, the viscosity of the synovial fluid is restored, which is a natural product of the body and Hyruan Inj. is an effective drug designed to replace the damaged synovial fluid found in osteoarthritis affected joints.

Composition

1 mL of Hyruan Inj. contains:
Active ingredient: Sodium Hyaluronate 10 mg
Excipients: Sodium chloride, Sodium phosphate dibasic, Sodium phosphate monobasic, Water for injection

Description

A viscosupplement for intra-articular use. It is a clear, colorless, odorless, and tasteless liquid. It is a natural product of the body and Hyruan Inj. is an effective drug designed to replace the damaged synovial fluid found in osteoarthritis affected joints.

Warnings and Precautions

1. Hyruan should not be administered to following patients:
Patients with a prior hypersensitivity to sodium hyaluronate.
2. Hyruan should be administered with caution to following patients:
1) Patients with a prior hypersensitivity to other drugs.
2) Patients with liver, kidney or heart failure, if it is observed that AET have occurred in the patients with prior history of these diseases.
3) Patients with skin disease or infection at injection site to be avoided.
3. Adverse effects:
1) Serious adverse effects:
Observe symptoms of shock (fainting, lightheadedness, rapid pulse, etc.) after administration should be made. In case observed on any noticed, administration should be discontinued and proper measures should be taken.
2) Other adverse effects:
1) Hypersensitivity: Erythema, edema, pruritus, urticaria, rash, angioedema, anaphylaxis, and so on may occur. In case these symptoms occur, administration should be discontinued and proper measures should be taken.
2) Arthralgia and myalgia: Occasionally, pain, swelling, stiffness, and so on may occur after injection, and these symptoms may occur.
3) Other adverse effects: Nausea, vomiting, and fever may occur.
4. General precautions:
1) It is strongly recommended that Hyruan should be given only after the removal of infection or caused by degenerative osteoarthritis of the joint when administered into severely inflamed joints. Hyruan Inj. is not effective for infection.
2) Since local pain can occasionally occur after administration of Hyruan Inj., should be advised patients that these symptoms are mild and self-limiting.
3) Since pain can be caused by the leakage of Hyruan Inj. or infection, it should be administered repeatedly and strictly under.